

ARTIN'S L -FUNCTIONS AFTER WEIL, GROTHENDIECK

BERNDT E. SCHWERTDFEGER

To Jo on his 3Fth birthday.

PREFACE

This paper summarizes some notes taken when comparing WEIL's presentation of ARTIN'S L -functions in [4] with GROTHENDIECK'S in [1].

Berlin, 15 November 2011

© 2011 Berndt E. Schwerdtfeger

v1.0

1. ARTIN L -FUNCTIONS ACCORDING TO WEIL

1.1. **DIRICHLET-series.** In general we will use the notation of WEIL in [5, VII] and [4, II]. We recap some basic notation.

Let k denote a global field, i.e. either an algebraic number-field (a finite field extension k/\mathbf{Q}) or an algebraic function-field of dimension 1 over a finite field of constants. For each place v of k its completion is denoted k_v . For finite places the local ring of integers is \mathfrak{o}_v and its maximal ideal \mathfrak{p}_v .

The free abelian group of all \mathfrak{p}_v is called group of *divisors* and is denoted \mathfrak{M} ; it is written *multiplicatively*. A divisor $\mathfrak{m} \in \mathfrak{M}$ is *positive* if the exponents of all \mathfrak{p}_v are positive. The positive divisors form the semigroup \mathfrak{M}_+ . In characteristic 0 (number-fields) \mathfrak{M} corresponds to the group of fractional ideals of k and \mathfrak{M}_+ to the ideals $\neq 0$ in the ring of integers \mathfrak{o}_k of k . In characteristic > 1 (function-fields) \mathfrak{M} corresponds to the *line bundles* on the associated complete curve – this aspect will not be delved into in this article. As we want to compare WEIL'S treatment of ARTIN'S L -functions with GROTHENDIECK'S we are particularly interested in the *function-field* case.

Let $\Omega_k = \text{Hom}(\mathbf{A}_k^\times / k^\times, \mathbf{C}^\times)$ be the group of (not necessarily unitary) characters on the idele class group of k . WEIL attaches to a function $c : \mathfrak{M}_+ \rightarrow \mathbf{C}$ an *extended DIRICHLET-series* belonging to k by

$$L(c, \omega) = \sum_{\mathfrak{m} \in \mathfrak{M}_+} c(\mathfrak{m}) \omega(\mathfrak{m})$$

This series is absolutely convergent for $\text{Re } \omega > \alpha + 1$ if and only if $|c(\mathfrak{m})| \leq C|\mathfrak{m}|^{-\alpha}$ for some $C > 0$ and some $\alpha \in \mathbf{R}$.

The coefficients $c(\mathfrak{m})$ are uniquely determined by the function $L(c, \omega)$.

2010 *Mathematics Subject Classification.* Primary 11M41; Secondary 11R42.

Key words and phrases. Artin-, Hecke L -functions, Dirichlet-series, Weil groups, Galois representations, inertia, decomposition group.

1.2. ARTIN-HECKE L -series. Let W_k be the WEIL group of k . The quotient by its derived group W_k/W'_k is canonically isomorphic to the idele-class group $\mathbf{A}_k^\times/k^\times$ by class field theory, let $\alpha_k : \mathbf{A}_k^\times/k^\times \xrightarrow{\sim} W_k/W'_k$ denote the ARTIN reciprocity map. So, we can also identify the characters $\omega \in \Omega_k$ on the idele-class group with the abelian characters $\chi : W_k/W'_k \rightarrow \mathbf{C}^\times$ (representations of degree 1).

Let X_k be the ring of representations of W_k (the WEIL group). If k'/k is a separable extension of degree d , then $W_{k'}$ is of index d in W_k and we can induce representations from $W_{k'}$ to W_k , denoted $\chi' \mapsto [\chi'; k'/k]$, and considered as a mapping $X_{k'} \rightarrow X_k$.

To each finite separable extension k'/k and to every character $\chi' \in X_{k'}$ there is attached a divisor $f_{k'}(\chi')$ of k' and an extended DIRICHLET series $L_{k'}(\chi')$ satisfying the following properties ([4, §73])

- (1) For $\deg \chi' = 1$ the divisor $f_{k'}(\chi') = f(\omega')$ is the conductor of $\omega' = \chi' \circ \alpha_{k'}$ and the L -series is the usual HECKE series to the Größencharakter ω' as in WEIL [5, VII] $L_{k'}(\chi') = L(\omega')$.
- (2) The mappings $\chi' \mapsto f_{k'}(\chi')$ and $\chi' \mapsto L_{k'}(\chi')$ are *homomorphisms* of the additive group $X_{k'}$ into the multiplicative groups of the divisors $\mathfrak{M}_{k'}$ of k' , and of the DIRICHLET series belonging to k' with initial coefficient 1, respectively.
- (3) For $k''/k'/k$ let $D(k''/k')$ be the *discriminant* of k'' over k' , let χ'' be any character of $W_{k''}$ of degree $n = \deg \chi''$. Then

$$\begin{aligned} f_{k'}([\chi''; k''/k']) &= N_{k''/k'}(f_{k''}(\chi''))D(k''/k')^n \\ L_{k'}([\chi''; k''/k']) &= [L_{k''}(\chi''); k''/k'] \end{aligned}$$

From these properties it is clear that these conditions determine the L -series for all characters.

For function fields WEIL has proven the ARTIN conjecture that these L -functions are *holomorph* for positive characters $\chi \neq 1$ ([2, II^e partie, §V, n^o 27-28]).

2. ARTIN L -FUNCTIONS ACCORDING TO GROTHENDIECK

2.1. Definition of Z - and L -functions. In [1], following WEIL [3, p. 507], GROTHENDIECK discussed the Z - and L -functions for a scheme Y of finite type over a finite field \mathbf{F}_q . Let the finite group G operate on the right such that $X = Y/G$ exists, let $\rho : G \rightarrow GL(V)$ be a representation in a finite dimensional vector space over a field F of characteristic 0 (classically $F = \mathbf{C}$, but we might also take $F = \mathbf{Q}_\ell$ or $F = \mathbf{C}_\ell$).

The ARTIN L -function is defined by

$$L(Y, G, X, t) = L(Y/X, \rho, t) = \prod_{x \in X^0} \frac{1}{\det(1 - \rho^{\natural}(x) t^{d(x)})}$$

where $X^0 =$ set of *closed* points of X , $d(x) = \deg x = [\kappa(x) : \mathbf{F}_q]$ is the residual degree and

$$\rho^{\natural}(x) = \frac{1}{|T_y|} \sum_{\substack{s \in G_y \\ s \mapsto \varphi_y}} \rho(s) \in \text{End}(V)$$

where $y \rightarrow x$ for a point y in the fiber Y_x , with G_y the *decomposition* group, T_y is the *inertia* group ($e(y/x) = |T_y| = e(x)$ is independent of y/x) and $\varphi_y \in G_y/T_y \simeq G(\kappa(y)/\kappa(x))$ is the FROBENIUS in the GALOIS group of the residual extension.

We have

$$\log L(Y/X, \rho, t) = \sum_{v \geq 1} c_v(Y, G, \rho) t^v / v$$

$$c_v(Y, G, \rho) = \sum_{x \in X(\mathbf{F}_{q^v})} \text{Tr}_\rho^{\natural}(\varphi_x)$$

where

$$\text{Tr}_\rho^{\natural}(\varphi_x) = \frac{1}{|T_y|} \sum_{\substack{s \in G_y \\ s \rightarrow \varphi_x}} \text{Tr}_\rho(s)$$

where $y \in Y(\mathbf{F}_{q^v})$ is above the point $x \in X(\mathbf{F}_{q^v})$ and φ_x is the FROBENIUS.

REFERENCES

- [1] Alexander Grothendieck, *Formule de Lefschetz et rationalité des fonctions L*, Sémin. Bourbaki **279** (1964/65), 1–15.
- [2] André Weil, *Courbes algébriques et variétés abéliennes*, Hermann, Paris, 1948. réimp. 1971.
- [3] ———, *Numbers of solutions of equations in finite fields*, Bull. Am. Math. Soc. **55** (1949), 497–508, available at <http://berndt-schwerdtfeger.de/wp-content/uploads/pdf/nf.pdf>.
- [4] ———, *Dirichlet Series and Automorphic Forms*, Lecture Notes in Math., vol. 189, Springer, Berlin, 1971.
- [5] ———, *Basic Number Theory*, Classics in Mathematics, Springer, Berlin, Heidelberg, New York, 1973, 1995.